


In-Furrow Use Guide for Cotton

Benefits of Priaxor® Fungicide

- Effective control of key soilborne diseases
- Improves yield and minimizes risk
- Provides Plant Health benefits for cotton


Priaxor Fungicide In-Furrow Provides Excellent Disease Control




2014-2015 Rhizoctonia challenged. Quitman, GA (n=2). Priaxor fungicide 4 fl oz/A.

Priaxor fungicide helps growers get the most out of every acre.

Rhizoctonia Root Rot in Cotton



Priaxor Fungicide: In-Furrow on Cotton for Rhizoctonia Seedling Disease



July 20, 2015 – Rhizoctonia challenged trial – Lubbock, TX. Rate per 1000 row/feet = 0.3 fl oz (Priaxor Fungicide)

Factors that Improve the Odds of Success and Yield Increases with Priaxor® Fungicide In-Furrow on Cotton

Soilborne diseases of cotton are prevalent in cotton fields throughout the U.S. Pythium, Fusarium, and Rhizoctonia seedling diseases are most common and cause significant yield losses of up to 5%. Recent university field trials have shown Priaxor fungicide used in-furrow can reduce the incidence and severity of these diseases and others while improving yield.

If any of these factors below are present on your farm, in-furrow applications of Priaxor Fungicide may be beneficial

- Fields with a history of:
 - Poor drainage
 - Slow establishment
 - Disease history (Rhizoctonia, Fusarium, Pythium)
- Early planting/cooler soils (<65° F)
- Stressful early season conditions (ie. sandblasting, etc)
- No-till/minimum till sites
- Low vigor varieties
- Seeding rate (<3 – 4 seeds/ft of row)


Best Use Recommendations

- Use Rate: 2.0 – 4.0 fl oz/A
- Application Volume: 2.5 – 10 GPA
- Conduct compatibility jar test before mixing with additional crop protection products
- Only use water as carrier (Do not use starter fertilizers as carrier)

Target Diseases

- Rhizoctonia seed, seedling, and root rot
- Fusarium seed rot and seedling blight*
- Pythium damping off*

*Suppression only.



Rate Per 1000 Row Feet (fl oz product)	Product Rate (fl ozs/A) row width (inches)									
	7" rows	15" rows	20" rows	22" rows	30" rows	32" rows	34" rows	36" rows	38" rows	40" rows
0.1	7.5	3.5	2.6	2.4	–	–	–	–	–	–
0.2	–	7.0	5.2	4.8	3.5	3.3	3.1	2.9	2.7	2.6
0.3	–	–	7.8	7.1	5.2	4.9	4.6	4.4	4.1	3.9
0.4	–	–	–	–	7.0	6.5	6.2	5.8	5.5	5.2
0.5	–	–	–	–	–	–	7.7	7.3	6.9	6.5
0.6	–	–	–	–	–	–	–	–	–	7.8

Do not exceed 8 fl oz/A. Refer to the label for other application directions.