

Merivon®

Xemium® Brand Fungicide


Summer Disease Control in Almonds

Benefits of Merivon® Fungicide

- Longer-lasting disease protection
- More consistent performance for maximum crop potential
- Advanced Plant Health benefits

Merivon fungicide provides more consistent performance for maximum disease control and crop potential.


Alternaria Leaf Spot Control with Merivon Fungicide



2012 J. Adaskaveg—University of CA—Riverside—Variety = Caramel
Treatments applied on 5/22 and 6/13 in 100 GPA. Disease evaluations made on 8/15.
For Alternaria disease incidence on leaves, 30-40 leaves from each of the three single-tree replications were evaluated for the presence of disease. Treatments are a subset of those evaluated in the trial which was sponsored by BASF and other product makers.



Almond Hull Rot with Merivon® Fungicide 2013



2013 Dr. Jim Adaskaveg—University of CA—Riverside—Treatments applied on 6/26, and 7/16 in 100 GPA. The first application was applied after a rain event to manage Monilinia hull rot. The second was at early suture opening to manage Rhizopus hull rot. Evaluations for disease were done on 8/16. The incidence of hull rot twig strikes was assessed based on evaluation of 150 spurs on each of four single-tree replications. Treatments are a subset of those evaluated in the trial which was sponsored by BASF and other product makers.

Best Use Recommendations

- Use Rate: 5.5–6.5 oz/A
- Optimal timing is 5.5 oz/A at full bloom
- Second application for summer diseases and/or Alternaria
- Merivon fungicide may be mixed with other products, such as insecticides
- Adjuvant flexible

Target Diseases

- Alternaria
- Anthracnose
- Botrytis blossom and shoot blight
- Brown rot/Blossom blight
- Green fruit/Jacket rot
- Hull rot suppression
- Scab
- Shothole
- Rust

Merivon Fungicide Scab Control



2013 Ag. Advisors Live Oak, CA Applications 1st = PF (3/15/13), 2nd = PF + 5 wks (4/19/13) and 3rd = PF + 8 wks (5/10/13). Photos 74 days after 3rd spray (7/23/13). Trial sponsored by BASF.

Merivon Fungicide Use Recommendation

Bloom			Spring		Summer	
Pink bud	Full bloom	Petal fall	2 weeks	5 weeks	May	June
Brown rot and Advanced Plant Health Merivon fungicide @ 5.5 oz/A			Scab, shothole, anthracnose and rust		Alternaria	



Always read and follow label directions.

Merivon, Pristine, and Xemium are registered trademarks of BASF. Quadris Top and Inspire are registered trademarks of Syngenta. Luna is a registered trademark of Bayer. Dyne-Amic is a registered trademark of CHEMTREC. PhD is a registered trademark of Arysta Life Science. Quash is a registered trademark of Valent.

©2015 BASF Corporation. All Rights Reserved. APN# 1403009 Merivon-Almonds-Summer-Diseases