

Chemical names of active ingredient

5,7,8-trimethyltolcol or RRR-alpha-tocopherol or D-alpha-tocopherol
5,8-dimethyltolcol or RRR-beta-tocopherol or D-beta-tocopherol
7,8-dimethyltolcol or RRR-gamma-tocopherol or D-gamma-tocopherol
8-methyltolcol or RRR-delta-tocopherol or D-delta-tocopherol

EU: Tocopherol-rich extract (E 306)

US: Natural mixed tocopherols

CAS-No.	D-alpha-tocopherol	59-02-9
	D-beta-tocopherol	16698-35-4
	D-gamma-tocopherol	54-28-4
	D-delta-tocopherol	119-13-1

EINECS-No.	D-alpha-tocopherol	200-412-2
	D-beta-tocopherol	240-747-1
	D-gamma-tocopherol	200-201-5
	D-delta-tocopherol	204-299-0

PRD-No.
30533885*

* The product is kosher

Articles*

50210253	10 kg plastic pail
50230963	190 kg metal drum

* packed under inert gas


Country of origin

USA

Description

Covi-ox® T-50 EU is a clear brownish-red viscous oil with mild taste and odor. It contains natural mixed tocopherols which are obtained from edible vegetable oils by suitable physical means. Covi-ox® T-50 EU is intended for use as an antioxidant in the dietary supplement and food industry.


D-alpha-tocopherol



$C_{29}H_{50}O_2$

Molar mass 430.7 g/mol


D-beta-tocopherol



$C_{28}H_{48}O_2$

Molar mass 416.7 g/mol


D-gamma-tocopherol



$C_{28}H_{48}O_2$

Molar mass 416.7 g/mol

D-delta-tocopherol



$C_{27}H_{46}O_2$

Molar mass 402.6 g/mol

Composition

Ingredients in descending order of weight:
tocopherol-rich extract* (E 306), sunflower oil.

* derived from soy beans

The typical tocopherol composition is:

14% D-alpha-tocopherol
2% D-beta-tocopherol
60% D-gamma-tocopherol
24% D-delta-tocopherol

Due to the variability of vegetable oil feedstocks used to derive the tocopherols, slight variations from this typical composition will be found, wide variations are not expected but they may occur.

Solubility

Insoluble in water, sparingly soluble in ethanol, miscible with organic solvents, vegetable oils and fats.

Specification

Assay (total tocopherols)	min. 500 mg/g
Assay (non-alpha-tocopherols)	min. 400 mg/g

For further information see separate document: "Standard Specification" (not for regulatory purposes) available via BASF's WorldAccount: <https://worldaccount.basf.com> (registered access).

Monographs

The active ingredient of Covi-ox® T-50 EU meets the monograph requirements of the current USP-NF and FCC (RRR-Tocopherols Concentrate, Mixed, Low-alpha-Type). Furthermore, the active ingredient complies with the current specification for food additives of EU Regulation (E 306).

Regulations

Covi-ox® T-50 EU is derived from IP-certified, not genetically modified raw materials, and is thus not subject to labelling under EU Regulations (EC) 1829/2003 and 1830/2003. Mixed tocopherols are approved for a use as an antioxidant and as a vitamin E source in most countries. However, specific regulations on the product in the respective countries and for its intended use have to be observed.

Stability, Storage and Handling

Stored in its unopened original packaging at ambient temperature max. 25 to 30 °C, the product is stable for at least 36 months. The product is sensitive to atmospheric oxygen, light and heat. It should therefore be stored in the tightly sealed, lightproof packaging in a cool place. Once opened, it is recommended to flush the packaging with an inert gas and to use the remaining contents as quickly as possible.

Applications

Dietary supplements, food and beverage products:

Covi-ox® T-50 EU is used as a natural antioxidant in a wide variety of food products and in industrial applications. The product has low volatility, excellent carry-through effect and is completely miscible with oils and fats. Covi-ox® T-50 EU should be added as early in the production process as possible. The amount of Covi-ox® T-50 EU to be used will vary with each particular application. For most oils and fats, the recommended concentration of tocopherols in the product will range from 100 ppm to 400 ppm relative to the fat or oil content. For oils containing highly polyunsaturated fatty acids, tocopherol concentrations of up to 2000 ppm relative to the oil content may be necessary. It is recommended to test 3 different mixed tocopherol concentrations to determine the optimum level which gives the desired antioxidant protection for each individual product.

Mixed tocopherols tend to darken over time, therefore it is recommended that in capsule applications, an amber shell is used.

Note

Covi-ox® T-50 EU must be handled in accordance with the Safety Data Sheet with the Safety Data Sheet.

This document, or any answers or information provided herein by BASF, does not constitute a legally binding obligation of BASF. While the descriptions, designs, data and information contained herein are presented in good faith and believed to be accurate, it is provided for your guidance only. Because many factors may affect processing or application/use, we recommend that you make tests to determine the suitability of a product for your particular purpose prior to use. It does not relieve our customers from the obligation to perform a full inspection of the products upon delivery or any other obligation.

NO WARRANTIES OF ANY KIND, EITHER EXPRESS OR IMPLIED, INCLUDING WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, ARE MADE REGARDING PRODUCTS DESCRIBED OR DESIGNS, DATA OR INFORMATION SET FORTH, OR THAT THE PRODUCTS, DESIGNS, DATA OR INFORMATION MAY BE USED WITHOUT INFRINGING THE INTELLECTUAL PROPERTY RIGHTS OF OTHERS. IN NO CASE SHALL THE DESCRIPTIONS, INFORMATION, DATA OR DESIGNS PROVIDED BE CONSIDERED A PART OF OUR TERMS AND CONDITIONS OF SALE.

July 2014